Consent for Nuclear Dobutamine Myocardial Perfusion

[,	authorize Dr. Reyes and his assist	tants to administer and conduct a
Dobutamine myocardial perfusion stress clinically significant heart disease; to evertimess for work or sports.		
I understand that I will be given a medic	cation named Dobutamine over a	neriod of approximately twelve
minutes through an intravenous site (IV injected through this same IV site once having the stress test, my electrocardiog will continue until I have completed the assistants consider significant, whichever). In addition, a radiopharmaceution my heart rate has reached a predet gram (EKG) and blood pressure we entire test, or I develop any abnormal management.	cal (Tc ^{99m} Myoview or Tl ²⁰¹⁾ will be termined target heart rate. While fill be continually monitored. The test
Every effort will be made to conduct the understand that just as with other types with a Nuclear Dobutamine Myocardi fainting, chest discomfort, leg cramps at effects associated with the radiopharmatequipped for such situations and that its necessary. I voluntarily accept the risks	of diagnostic tests there are potent ial Perfusion stress test. These ind nd very rarely heart attacks or sud ceutical injection. I further unders professional personnel are trained	tial side effects and risks associated clude episodes of lightheadedness, den death. There are no risks or side stand that the clinic is properly d to administer any emergency care
Patient Signature		Date
Witness Signature		Date